

Marie Atallah, PhD
Sutter Health
doc@marieatallah.com

Abstract

Ileostomy, a surgical intervention required for conditions such as inflammatory bowel disease (IBD), colorectal cancer, trauma, and sepsis, can be a transformative intervention and imposes substantial physical and psychological challenges on patients. While the physical aspects of ileostomy care are well addressed, the psychological burden — marked by anxiety, depression, and diminished quality of life — remains under-recognized in clinical practice. This perspective article underscores the critical role of mental health in ileostomy recovery, highlighting variations in psychological distress and recovery trajectories across adult patient populations. For some, such as those with IBD, the ileostomy represents relief from debilitating symptoms. For others, including trauma or cancer patients, the stoma may symbolize loss, exacerbating emotional distress. Unaddressed psychological distress can impede recovery through physiological mechanisms (e.g., dysregulated inflammation, impaired immunity) and behavioral pathways (e.g., non-adherence to care, social withdrawal). Compounding this issue is a pervasive gap in clinician training: few healthcare providers are equipped to address the unique mental-health needs of ostomy patients. This article synthesizes emerging evidence to advocate for integration of mental-health support into ileostomy care, emphasizing its potential to optimize outcomes across diverse adult populations and drawing parallels to successful mental-health integration in other chronic and acute conditions.

Variable Psychological Impact of Ileostomy: Context Matters

The psychological burden of an ileostomy is not monolithic — it is shaped by the patient’s underlying condition and the narrative surrounding their journey to surgery.¹ For some, such as IBD patients, the ileostomy represents relief after years of suffering.^{2 3} For others, like trauma or cancer patients, it has a more sudden occurrence and may become an unwelcome reminder of a life-altering event.^{4 5}

IBD Patients: Liberation from Chronic Suffering

For many IBD patients, ileostomy offers respite from years of debilitating symptoms, including pain, malnutrition, and diarrhea. Unsurprisingly, 70-80% of IBD patients report significant improvements in quality of life post-ileostomy, with many describing the stoma as “a second chance at life.”^{2 3} Even among patients who experience improved quality of life, challenges such as body-image disturbance, fear of leaks, and anxiety about intimacy persist.^{3 4} Notably, younger IBD patients report higher rates of depression post-surgery, potentially due to the unique challenges of their life stage—such as navigating relationships, careers, and social dynamics while adapting to life with a stoma.^{4 5}

Cancer Patients: A Symbol of Illness and Fear

For cancer patients, an ileostomy often represents a tangible reminder of their diagnosis and treatment. The stoma may compound the psychological toll of cancer, evoking fears of recurrence, social rejection, or a diminished sense of self. Colorectal-cancer patients with stomas report clinically significant anxiety, frequently linked to stoma-related concerns such as odor, leakage, or visibility of the pouch.^{5 6} Unlike IBD patients—who may share a narrative of the stoma as a source of relief—cancer patients often struggle to reconcile the stoma with their identity, leading to prolonged emotional distress.⁶

Trauma Patients: An Unplanned Life-Altering Event

Trauma patients—such as those undergoing ileostomy following accidents, acute medical emergencies, or surgical complications—face a distinct set of challenges. For these individuals, the stoma is often the result of an unplanned, life-altering event, leaving little time for psychological preparation.^{7 8} They may struggle with feelings of vulnerability or loss of control, and the stoma becomes a physical manifestation of their traumatic experience. Unlike cancer patients, who may have some time to mentally prepare for the possibility of a stoma, trauma patients are thrust into this reality abruptly, often without the opportunity to process its implications.⁸




Figure 1. The Variable Psychological Impact of Ileostomy: Context Matters illustrates the overlapping and distinct psychological challenges faced by ileostomy patients based on their underlying conditions: inflammatory bowel disease (IBD), trauma, and cancer. While all groups share universal challenges, each population also faces unique concerns shaped by their journey to an ileostomy. Tailored interventions are essential to address these nuanced needs.

How Distress Undermines Recovery: Mechanisms and Evidence

Psychological distress is not merely a subjective experience—it exerts tangible effects on postoperative healing through physiological and behavioral pathways.^{9–15}

Physiological Pathways: Stress, Inflammation, and Immunity

Chronic stress activates the hypothalamic–pituitary–adrenal (HPA) axis, triggering systemic release of cortisol and pro-inflammatory cytokines. Elevated cortisol levels impair fibroblast proliferation and collagen synthesis, **can** delay wound healing and increasing the risk of surgical-site infections.⁹ Concurrently, dysregulated inflammation exacerbates underlying conditions—for example, in IBD patients, psychological distress has been linked to significantly higher odds of disease flares.¹⁰ Emerging research also implicates stress-induced gut dysbiosis in stoma-related complications. Animal models demonstrate that chronic stress alters gut microbiota composition, reducing microbial diversity and impairing mucosal immunity—a finding with direct relevance to ileostomy patients.^{10–11} For these patients, the diverted digestive system and absence of the colon create a unique vulnerability to microbiome disturbances. Stress-induced changes in gut flora **can** exacerbate complications such as small intestinal bacterial overgrowth (SIBO), leakage, and peristomal skin irritation.¹¹

Behavioral Pathways: Nonadherence, Avoidance, and Isolation

Distressed ileostomy patients are more likely to engage in behaviors that jeopardize recovery, including nonadherence to stoma care, intentional fluid restriction, and social withdrawal.¹² Anxiety about pouch changes, fear of high output, or embarrassment about the stoma can lead to complications such as skin breakdown, dehydration, and further isolation.^{12–13} These poor outcomes, in turn, exacerbate distress, creating a feedback loop that prolongs recovery, extends hospital stays, and escalates healthcare costs.^{13–15}



Figure 2. The Cycle of Distress and Complications in Ileostomy Patients.

This figure illustrates the self-reinforcing cycle in which psychological distress leads to nonadherence to stoma care, resulting in physical complications such as skin breakdown and dehydration. These complications, in turn, exacerbate psychological distress, prolong recovery, extend hospital stays, and escalate healthcare costs. The cycle underscores the bidirectional relationship between mental health and physical outcomes in ileostomy patients.

Training Gap: Why Clinicians Are Ill-Equipped to Respond

Although surgeons frequently recognize the psychological challenges associated with ostomy care, formal distress screening and mental-health support remain inconsistently implemented in surgical and cancer-care settings.¹⁶ Even when distress is recognized, few mental-health professionals are trained to address stoma-specific challenges. Therapists often feel ill-equipped to manage fears of leakage, odor, or social rejection—issues central to the psychological burden of ileostomy patients.^{16 17} Without targeted training, mental-health providers may struggle to contextualize patients' concerns, further perpetuating the care gap. Bridging this gap requires interdisciplinary collaboration among surgeons, stoma nurses, and mental-health professionals.^{16 17}

Call to Action

Addressing mental health in ileostomy care requires a multifaceted approach. Specialized training programs must target both surgical and mental-health teams. Mental-health modules should be integrated into ostomy-care certification curricula, and distress screening (e.g., PHQ-9 or OAI-23) embedded into pre- and post-operative care.^{18 19} Embedding these screenings into EHR systems can automate referrals for high-risk patients. Ostomy-specific mental-health interventions—such as tailored CBT and peer-support networks—have shown promise in improving outcomes and reducing complications.^{20 21} Finally, policy and reimbursement reform are needed to sustain these programs, along with research to address disparities and identify biomarkers of stress-related complications.⁸

Conclusion

The mind–body dichotomy is a false construct in ileostomy care. Psychological distress is not ancillary—it is a critical mediator of recovery. For IBD patients, the stoma may symbolize hope; for cancer survivors, despair. These divergent narratives demand individualized mental-health support to empower patients to navigate recovery with resilience and dignity.

References

1. Starčević M, et al. *Quality of life and psychological distress after stoma surgery: a systematic review. Colorectal Disease.* 2025.
2. Pineton de Chambrun G, et al. *Quality of life of patients with inflammatory bowel diseases who live with an ostomy: results of a French national survey. Journal of Crohn's and Colitis.* 2018;12(Suppl 1):S466. doi:10.1093/ecco-jcc/jjx180.828
3. Davis S, et al. *Impact of stoma on lifestyle and health-related quality of life in patients living with stoma. Indian Journal of Palliative Care.* 2020;26:84–90. doi:10.4103/IJPC.IJPC_111_19
4. Ananthakrishnan AN, et al. *Similar risk of depression and anxiety following surgery or hospitalization for Crohn's disease and ulcerative colitis. American Journal of Gastroenterology.* 2013;108(4):594–601. doi:10.1038/ajg.2012.471
5. Ye M, et al. *Supportive care needs and psychological distress among colorectal cancer patients with a stoma. Supportive Care in Cancer.* 2023;31:6031–6039.
6. Soelling S, et al. *Living with a stoma: a qualitative study of psychosocial adaptation. Journal of Wound, Ostomy and Continence Nursing.* 2025.
7. Clark J, et al. *Psychological preparation for surgery: systematic review. Patient Education and Counseling.* 2020;103(12):2453–2464.
8. Lederer A-K, et al. *Impact of surgery-related emotional distress on long-term outcome after colorectal surgery. Journal of Clinical Medicine.* 2024;13(18):6406. doi:10.3390/jcm13186406
9. Saito H, et al. *The effect of stress on fibroblast proliferation and collagen synthesis. Surgery Today.* 1997;27:518–522.
10. Li C, et al. *Psychological stress and inflammatory bowel disease outcomes: mechanisms and management. Inflammatory Bowel Diseases.* 2022;28(9):1331–1342.
11. Rao SSC, et al. *Brain fog and D-lactic acidosis in small intestinal bacterial overgrowth: case series. Clinical and Translational Gastroenterology.* 2018;9:e196.
12. Vonk-Klaassen SM, et al. *Ostomy-related problems and their impact on quality of life: a systematic review. Journal of Clinical Nursing.* 2016;25(11–12):1481–1502.
13. Taneja C, et al. *Clinical and economic burden of peristomal skin complications. Journal of Wound, Ostomy and Continence Nursing.* 2019;46(2):143–149.
14. DiMatteo MR, Lepper HS, Croghan TW. Depression is a risk factor for noncompliance with medical treatment: meta-analysis of the effects of anxiety and depression on patient adherence. *JAMA.* 2000; 284(20):2430–2435. doi:10.1001/jama.284.20.2430
15. Andersen RM, et al. *Cost-effectiveness of interventions preventing stoma complications. Expert Review of Pharmacoeconomics & Outcomes Research.* 2024;24(1):13–25.
16. Saunders CL, et al. *Training needs and priorities for stoma care professionals. BMJ Open.* 2023;13:e071234.
17. Polidano K, et al. *The experience of living with an ostomy in young adults: a qualitative meta-synthesis. Qualitative Health Research.* 2021;31(7):1262–1275.
18. Kroenke K, et al. *The PHQ-9: validity of a brief depression severity measure. Journal of General Internal Medicine.* 2001;16(9):606–613.

18. Simmons KL, Smith JA, Maekawa A. *Development and psychometric evaluation of the Ostomy Adjustment Inventory-23. Journal of Wound, Ostomy and Continence Nursing.* 2009;36(1):69–76. doi:10.1097/WON.0b013e3181919b7d
19. Glass A, Monroe M, Davies C. *The effect of a multidisciplinary stoma team on anxiety and distress in patients undergoing colorectal stoma surgery. Journal of Oncology Navigation & Survivorship.* 2022;13(7).
20. Lim SH, et al. *Ostomy-specific cognitive-behavioral intervention: pilot outcomes. Patient Education and Counseling.* 2019;102(8):1543–1549.